

Cosmic Ray Anisotropy Workshop 2023

Tuesday 16 May 2023 - Friday 19 May 2023

Loyola University - Chicago



CRA 2023

sixth Cosmic Ray Anisotropy Workshop

May 16-19, 2023
Loyola University - Chicago



Book of Abstracts

Contents

Registration	1
Welcome	1
Transport of Galactic Cosmic-Ray Nuclei	1
Cosmic rays in a turbulent interstellar medium: Recent progress and open questions	1
Indirect Measurements of galactic Cosmic Rays	1
Recent Progress in Direct Measurements of Cosmic Rays (remote)	1
CR anisotropy with IceCube	1
CR anisotropy with IceTop - Rasha Abbasi	2
Registration	2
Selected Aspects of Magnetized Turbulence (with implications for CRs)	2
Second order Fermi Acceleration in Galaxy Clusters	2
CR Acceleration in Turbulent Magnetic Reconnection (remote)	2
Cosmic ray diffusion in anisotropic MHD turbulence	2
Constraining the sources of ultra-high-energy cosmic rays with the Pierre Auger Observatory	2
Results from the High Altitude Water Cherenkov Observatory	2
New Models of the Magnetic Field of the Galaxy	3
UHECR anisotropies	3
Registration	3
Heliosphere in the Local Interstellar Medium	3
The IBEX Ribbon and its Relation to the Solar-Interstellar Interaction (remote)	3
Living in the Bubble: structure, formation, and how it affects us	3
Review of the Local ISM	3
Cosmic ray pitch angle anisotropies in the Very Local Interstellar Medium	3

Turbulence in the local ISM	4
Modeling of cosmic-ray anisotropy at TeV energies in an MHD model heliosphere	4
Heliosphere and TeV CR anisotropy	4
Configuration of Interstellar Magnetic Field Surrounding the Heliosphere (remote) . . .	4
Cross-correlation studies with UHECRs (remote)	4
Local Sources of Cosmic Rays	4
Wrapping up and Collecting thoughts on Cosmic Ray Astrophysics	4
Cosmic Ray Anisotropy with the Telescope Array	5
Welcome	5
The Relationship between the Local ISM Distribution and Milky Way Spiral Structure . .	5
Modeling of cosmic-ray anisotropy at TeV energies in an MHD model heliosphere	5
Review of the Local ISM	5
Results from TA	6
IceCube: The first Decade of Neutrino Astronomy	6
Cosmic rays in a turbulent interstellar medium: Recent progress and open questions . .	6
The IBEX Ribbon and its Relation to the Solar-Interstellar Interaction	6
The curious maximum-rigidity distribution of ultra-high-energy-cosmic-ray accelerators	7
Turbulence and CR propagation (remote)	7
Cosmic Ray Anisotropy with the Telescope Array	7
Observation of cosmic-ray anisotropy with LHAASO	8
Observation of cosmic-ray anisotropy with LHAASO (remote)	8
Recent Results in Cosmic-Ray Astrophysics	8

7

Registration

8

Welcome

Corresponding Author: rasha@icecube.wisc.edu

10

Transport of Galactic Cosmic-Ray Nuclei

11

Cosmic rays in a turbulent interstellar medium: Recent progress and open questions

12

Indirect Measurements of galactic Cosmic Rays

13

Recent Progress in Direct Measurements of Cosmic Rays (remote)

Author: Eun-Suk Seo¹

¹ *University of Maryland*

Corresponding Author: seo@umd.edu

14

CR anisotropy with IceCube

Corresponding Authors: rasha@icecube.wisc.edu, mcnelly_ft@mercer.edu

15

CR anisotropy with IceTop - Rasha Abbasi

16

Registration

17

Selected Aspects of Magnetized Turbulence (with implications for CRs)

18

Second order Fermi Acceleration in Galaxy Clusters

19

CR Acceleration in Turbulent Magnetic Reconnection (remote)

Author: Elisabete de Gouveia Dal Pino¹

¹ IAG - Universidade de São Paulo

Corresponding Author: dalpino@iag.usp.br

20

Cosmic ray diffusion in anisotropic MHD turbulence

21

Constraining the sources of ultra-high-energy cosmic rays with the Pierre Auger Observatory

22

Results from the High Altitude Water Cherenkov Observatory

23

New Models of the Magnetic Field of the Galaxy

24

UHECR anisotropies

25

Registration

26

Heliosphere in the Local Interstellar Medium

27

The IBEX Ribbon and its Relation to the Solar-Interstellar Interaction (remote)

28

Living in the Bubble: structure, formation, and how it affects us

29

Review of the Local ISM

30

Cosmic ray pitch angle anisotropies in the Very Local Interstellar Medium

31

Turbulence in the local ISM

32

Modeling of cosmic-ray anisotropy at TeV energies in an MHD model heliosphere

33

Heliosphere and TeV CR anisotropy

Author: Ming Zhang¹

¹ *Florida Institute of Technology*

Corresponding Author: mzhang@fit.edu

34

Configuration of Interstellar Magnetic Field Surrounding the Heliosphere (remote)

35

Cross-correlation studies with UHECRs (remote)

Author: Federico Urban^{None}

Corresponding Author: federico.urban@fzu.cz

36

Local Sources of Cosmic Rays

37

Wrapping up and Collecting thoughts on Cosmic Ray Astrophysics

38

Cosmic Ray Anisotropy with the Telescope Array

Cosmic ray anisotropy workshop / 39

Welcome

40

The Relationship between the Local ISM Distribution and Milky Way Spiral Structure

41

Modeling of cosmic-ray anisotropy at TeV energies in an MHD model heliosphere

Author: Takashi Sako^{None}**Co-author:** On behalf of the Tibet ASgamma Collaboration**Corresponding Author:** tsako@icrr.u-tokyo.ac.jp

Past and on-going cosmic-ray experiments have reported small ($\sim 0.1\%$) anisotropies in the arrival directions of TeV cosmic rays observed at the Earth.

We are attempting to estimate anisotropic features at the heliospheric boundary by applying the idea of Liouville mapping to the data of the Tibet AS γ experiment.

Our preliminary results have indicated small, possibly spurious, anisotropic structures, with angular scales of $\sim 10^\circ$ in the cosmic-ray intensity distribution at the heliospheric boundary, and we expect that the higher-order residues of these structures at the heliospheric boundary could be removed if the stochastic scattering of cosmic-ray particles by magnetic irregularities inside the heliosphere are somehow taken into account in the mapping process.

In this presentation we will present the latest results of our improved intensity-mapping method.

42

Review of the Local ISM

Authors: Jeffrey Linsky¹; Eberhard Moebius²

¹ University of Colorado

² *University of New Hampshire*

Corresponding Authors: eberhard.moebius@unh.edu, jlinsky@jila.colorado.edu

The local interstellar medium (LISM), consisting of neutral and ionized gas, dust, and magnetic fields, is the environment for the heliosphere and stellar astrospheres. LISM gas and magnetic fields penetrate deeply into these environments. The LISM consists of the partially ionized gas clouds extending to about 10 pc from the Sun, the surrounding Local Bubble with its fully ionized hydrogen gas extending to several hundred parsecs from the Sun, and interactions with the external Galaxy. Absorption in resonance lines, primarily in the ultraviolet, observed by the Hubble Space Telescope and in situ measurements by the Voyager, New Horizons, Ulysses, and IBEX spacecraft are providing important data concerning the composition and properties of the LISM. This talk will review what we have been learning about the LISM - its structures, thermal and non-thermal properties, turbulence, ionization, inhomogeneity, and pressure balance among its components. As the Sun speeds through the LISM at 25 km/s, large changes in the size and properties of the heliosphere occur when it passes through cold dense clouds, partially ionized gas, or fully ionized gas. Cosmic rays traverse the different structures in the LISM and are modified by these environments.

43

Results from TA

44

IceCube: The first Decade of Neutrino Astronomy

Author: Francis Halzen¹

¹ *University of Wisconsin–Madison*

Corresponding Author: halzen@icecube.wisc.edu

45

Cosmic rays in a turbulent interstellar medium: Recent progress and open questions

Author: Philipp Mertsch^{None}

Corresponding Author: pmertsch@physik.rwth-aachen.de

TBD

46

The IBEX Ribbon and its Relation to the Solar-Interstellar Interaction

Author: Eric Zirnstein¹

¹ *Princeton University*

Corresponding Author: ejz@princeton.edu

NASA's Interstellar Boundary Explorer (IBEX), an Earth-orbiting Small Explorer spacecraft, measures energetic neutral atom (ENAs) produced primarily by charge exchange in the outer heliosphere. There are two main sources of ENAs. The first, called the "globally distributed flux", is formed from neutralization of interstellar pickup ions that were preferentially accelerated at the heliospheric termination shock and advected with the bulk plasma flow through the inner heliosheath. Once these ions experience charge exchange, they may travel back towards in the inner heliosphere and be detected by IBEX. The second main source, called the "Ribbon", is a narrow enhancement of ENA emissions circling the sky that come from outside the heliopause. It is widely believed that the Ribbon is ordered by the draping of the local interstellar magnetic field (ISMF) around the heliosphere and has allowed us to pinpoint the pristine (far from the heliosphere) ISMF strength and orientation. I will present a review of Ribbon modeling studies that analyze the relationship between the Ribbon and the draped ISMF, the importance of understanding particle pitch angle scattering outside the heliopause, and the evolving solar wind that serves as the primary source of the Ribbon flux through the secondary ENA process.

47

The curious maximum-rigidity distribution of ultra-high-energy-cosmic-ray accelerators

Author: Foteini Oikonomou¹

Co-authors: Domenik Ehlert ; Michael Unger

¹ *Norwegian University of Science and Technology*

Corresponding Author: foteini.oikonomou@ntnu.no

A standard assumption among models of candidate source populations of ultra-high energy cosmic rays (UHECRs) is that all sources in a candidate source population accelerate particles to the same maximum energy. Motivated by the fact that candidate astrophysical accelerators exhibit a vast diversity in terms of their relevant properties, such as luminosity, Lorentz factor, and magnetic field strength, we study the compatibility of a population of sources with non-identical maximum cosmic-ray energies with the observed energy spectrum and composition of UHECRs at Earth. For this purpose, we compute the UHECR spectrum emerging from a population of sources with a power-law, or broken-power-law, distribution of maximum energies applicable to a broad range of astrophysical scenarios. We find that for a wide range of studied models, the maximum energies of the UHECR accelerators must be nearly identical in order to be compatible with the UHECR data, in stark contrast to the variance expected for the astrophysical source models considered. A substantial variance of the maximum energy is only consistent with the UHECR data if the maximum energies of the UHECR sources follow a broken power-law distribution with a very steep spectrum above the break. However, in this scenario, the individual source energy spectra must be unusually hard with increasing energy output as a function of energy. These findings have implications for the arrival-direction distribution of UHECRs.

48

Turbulence and CR propagation (remote)

49

Cosmic Ray Anisotropy with the Telescope Array

Authors: Jihyun Kim¹; for the Telescope Array Collaboration^{None}

¹ *University of Utah*

Corresponding Author: jihyunkim316@gmail.com

Ultra-high energy cosmic rays (UHECRs) are charged particles that are extremely energetic, with $E > 10^{18}$ eV. They impinge on the Earth's atmosphere from outer space. The Telescope Array experiment, the largest UHECR observatory in the northern hemisphere, is situated in the western desert of Utah, USA, and has been collecting data continuously since May 2008. It is designed to detect extensive air showers (EAS) which are a cascade of subatomic particles induced by a primary UHECR particle interacting with the atmosphere. The Telescope Array uses two types of detectors: fluorescence detectors (FDs) and scintillator surface detectors (SDs). The FDs measure the scintillation light produced when the shower travels through the atmosphere's gas while the SDs sample the EAS's footprint at ground level. To understand the origin of UHECRs, we analyze the distribution of their arrival directions, looking for indications of anisotropic patterns. In this presentation, we summarize the results of the anisotropy study of UHECR events observed by the Telescope Array SD array, taking full advantage of its high duty cycle.

50

Observation of cosmic-ray anisotropy with LHAASO

Authors: Wei Gao¹; David Ruffolo²; Wei Liu³; Jiayin He⁴

¹ *Institute of High Energy Physics*

² *Mahidol University*

³ *IHEP*

⁴ *Purple Mountain Observatory, CAS*

Corresponding Author: gaowei@ihep.ac.cn

Observation of cosmic-ray anisotropy with LHAASO

51

Observation of cosmic-ray anisotropy with LHAASO (remote)

52

Recent Results in Cosmic-Ray Astrophysics

Corresponding Author: carmelo.evoli@gssi.it