



# The IBEX Ribbon and its Relation to the Solar-Interstellar Interaction


Eric J. Zirnstein

Space Physics Group

Department of Astrophysical Sciences, Princeton University

Princeton, NJ 08540, USA

CRA 2023 Workshop, May 18, 2023




## *Topical questions:*


- What is the IBEX Ribbon and how is it formed?
- How is the Ribbon related to the Interstellar Medium?
- What are the macro and micro-physical processes governing the Ribbon's properties?
- What is the latest information we have about the local interstellar magnetic field (ISMF)?

- **Part 1: Ribbon intensity**
  - Small scales from PUI transport
  - Large scales from ISMF draping
  - All scales from neutral SW/PUI source
- **Part 2: Ribbon structure**
  - Both PUI dynamics and ISMF draping, turbulence
- **Part 3: Ribbon position and circularity**
  - \*Mostly\* due to ISMF draping
- **Part 4: Comparison with Voyager data**
  - Simulated ISMF independently derived from IBEX data compared to Voyager in situ field measurements
- **Part 5: Ribbon evolution**
  - Governed by solar cycle
- **Conclusions**

# What is the IBEX Ribbon?





# How is the Ribbon Formed?




- 1) Interstellar neutrals propagate into the heliosphere
- 2) Neutral charge exchanges with solar wind ion
- 3) Solar wind ion is neutralized (“primary ENA”), continues its original trajectory radially outwards and likely escapes the heliosphere
- 4) Neutral solar wind particle ionizes in interstellar medium, “PUI” gyrating in local field
- 5) PUI charge exchanges with another interstellar neutral, becoming neutralized (“secondary ENA”)
- 6) Some secondary ENAs propagate back inside the heliosphere
- 7) Some secondary ENAs are observed by IBEX

# How is the Ribbon Formed?



# Draping of the ISMF and its Effects on the Ribbon

Zirnstein et al. (2015, ApJL)




- Polar cut of Ribbon production rate
  - Isocontours of ribbon: 70%, 25%, 10%
- Surface representing  $\mathbf{B} \cdot \mathbf{r} = 0$  ( $\mathbf{r}$  is IBEX line of sight,  $\mathbf{B}$  is ISMF) is magenta curve
- Distance from heliopause at which 50% of Ribbon flux is accumulated is white curve
- Ribbon flux is concentrated near  $\mathbf{B} \cdot \mathbf{r} = 0$ , close to the heliopause
- $\mathbf{B} \cdot \mathbf{r} = 0$  surface is “warped” as a function of distance from the heliopause due to compression and draping of the ISMF


# 3D Ribbon Structure

Zirnstern et al. (2015, ApJL)

Ribbon Source (ESA 3)




Ribbon Source (ESA 3)




# 3D Ribbon Structure

Zirnstein et al. (2015, ApJL)

Ribbon Source (ESA 5)





Ribbon Source (ESA 5)





# 3D Ribbon Structure


Zirnstern et al. (2015, ApJL)



McComas et al. (2000, JGR)

# Draping of the ISMF and its Effects on the Ribbon


Zirnstein et al. (2020, ApJ)




# Ribbon Structure is Affected by Interstellar Turbulence

Zirnstain et al. (2020, ApJ)

- The power level and correlation scale of turbulence in the interstellar medium may significantly affect the Ribbon's shape and position in the sky





$\lambda_U = 10 \text{ au}$      $50 \text{ au}$      $100 \text{ au}$      $500 \text{ au}$




Differential Flux [ $\# / (\text{cm}^2 \text{ s sr keV})$ ],  $E = 1.11 \text{ keV}$

0    100    200    300    400



# Ribbon Position is Energy-dependent

Zirnstien et al. (2016, ApJL)




- Draping of the ISMF around the heliosphere creates spatially-changing Ribbon source
- Higher energy *secondary* (Ribbon) ENAs originate on average farther from the heliopause due to longer mean free path of *primary* ENAs
  - Causes Ribbon observed at higher energies (e.g., ESA 6) to appear to have a larger radius than at lower energies (e.g., ESA 2)
- Ribbon sources still overlap



# Ribbon Energy-dependent Circularity

Zirnstern et al. (2016, A&A)

## Ribbon centered frame





## $B \cdot r$ angle




# Ribbon Energy-dependent Circularity

Zirnstien et al. (2016, ApJL)





- Modeling the IBEX Ribbon at each ESA with different ISMF configurations (magnitude and direction) allows us to derive the most likely *pristine* ISMF outside our heliosphere



# Comparison with Voyager Observations of ISMF


Rankin et al. (2023, ApJL)




# Comparison with Voyager Observations of ISMF

Rankin et al. (2023, ApJL)


## Raw Voyager Data & Model



## Substitute Voyager $B_r$ from Model



Distance from Sun [au]






# Ribbon Evolution

Zirnstern et al. (2023, in press)


- Time-dependent results shown for ESA 3 from 2009-2014
  - *Again, spatial retention model is scaled up by factor of 4*
- Overall, there is a global decrease in ENA intensity at similar rates seen in the data and models
- Not much to distinguish between models



# Ribbon Evolution

Zirnstern et al. (2023, in press)


- Time-dependent results shown for ESA 3 from 2015-2019
  - Again, spatial retention model is scaled up by factor of 4
- Fluxes decrease until ~2016, then start increasing in southern hemisphere in ~2019
- Again, not much to distinguish between models



# Evolution of Ribbon Radii Over Time

Zirnstern et al. (2023, to be submitted)

Comparison of Ribbon Radius Over Time




- Weak scattering results consistently overestimate ribbon radius (not new result; Zirnstern et al. 2021)
- Strong scattering results consistently match IBEX ribbon radius, though data uncertainties likely underestimated at ESA 5-6

# Energy-averaged Centers and Radii

Zirnstien et al. (2023, to be submitted)

- Both data and model ribbon centers, averaged over energy, lie close to each other on the  $B-V$  plane
  - NOTE: We did not include ESA 6 in the average due to missing data and high systematic uncertainty unaccounted for**
  - Slight offset from data likely due to a few degree offset of the simulated interstellar magnetic field direction
- Ribbon centers do not appear to significantly change over time except from 2016 to 2019
- Ribbon radii do not appear to significantly change over time (in either model or data)
- Weak scattering radius overestimates data b/c ENA source is distributed much farther from heliosphere, reflecting closer to ‘great circle’
- Strong scattering radius matches data within uncertainties, with not obvious trend over time

## Comparison of Energy-averaged Ribbon Properties Over Time




# Conclusions

- Ribbon shape, position, and intensity is strongly influenced by:
  - the draping of the ISMF around the heliosphere
  - the neutral SW/PUI distribution as a function of latitude and time
  - Small-scale PUI dynamics outside the heliopause
- Multiple “observables” are available to try to differentiate different Ribbon source models
  - Intensity, cross-section shape and width, position in the sky, circularity (radius, center location)
- Modeling ribbon to fit to IBEX observations allowed us to derive best-fit ISMF vector far from heliosphere
  - The results independently compare well to Voyager data, understanding that the measured  $B_r$  component has significant systematic uncertainties
- Ribbon evolution over the solar cycle is replicated well in both weak and spatial retention models, making them essentially indistinguishable
  - Largest difference between models is the low intensity of the spatial retention model, requiring at least factor of 4 increase to match data
- Ribbon centers and radii, however, do reveal noticeable differences between the models
- Based on our results, the spatial retention model is favored in reproducing the data as a function of time and geometry
- However, modifications to the spatial retention theory is required to fix the intensity problem
  - Scattering effects near  $B \cdot r = 0$  amplifying distribution with pitch angles closer to  $90^\circ$  would increase ribbon flux at 1 au (Zirnstein et al. 2020)


# Extra Slides

# Evolution over Sections of Sky

Zirnstien et al. (2023, in press)




- We show results for **data**, **spatial retention model**, and **weak scattering model**
- Also show results where we scale IPS SW speeds to match OMNI (solid curves), and don't scale to OMNI (dashed curves)
- Numbers next to labels show factors used to scale the data/results such that the time-average equals 1



# Evolution of Ribbon Centers Over Time

Zirnstern et al. (2023, in press)


Comparison of Ribbon Center Over Time



- While not overlapping exactly, the general trend in energy/latitude of ribbon centers is reproduced in the models
- The spatial retention model compares slightly better, particularly in 2014-2015 – just a systematic offset





# Ribbon Source in “Weak Scattering”





# Distance to the Ribbon Source


Spatial Retention ( $1/\tau_s \ll 1/\tau_x$ )




Spatial Retention ( $1/\tau_s \gg 1/\tau_x$ )




Differential Flux [ $\#/(cm^2 s sr keV)$ ],  $E = 1.11 keV$



Distance to 50% of Total Flux [au],  $E = 1.11 keV$



# Ribbon Cross-section Shape



- “Spatial retention” produces shape skewed away from Ribbon center (leftward)
- “Isotropic scattering” produces shape skewed towards Ribbon center (rightward)
- “Weak scattering” produces the most symmetric shape

