

A measurement of the cosmic ray anisotropy at and above 10^{14} eV

Friday, 28 October 2011 15:00 (20 minutes)

The EAS-TOP Extensive Air Shower array was located at Campo Imperatore (2005 m a.s.l., latitude $42^{\circ}27'N$, longitude $13^{\circ}34'E$, INFN Gran Sasso National Laboratory). It took cosmic ray data in the energy range 10^{13} eV- 10^{16} eV from the end of 1980s up to 2000. A first data-set (including 4 years of data) was exploited for the measurement of the cosmic ray anisotropy at $E \approx 10^{14}$ eV (Ap. J. 470, 1996, 501). At this energy, the EAS-TOP results demonstrated that the main features of the anisotropy (i.e., of cosmic-ray propagation) are similar to those measured at lower energies (10^{11} - 10^{14} eV), both with respect to amplitude ($(3-6) \sim 10^{-4}$) and phase ($(0-4)$ hr local sidereal time (LST)). Thanks to the final data-set (spanning over 8 years) the EAS-TOP measurement could be extended to higher energy, about 4×10^{14} eV. The observed anisotropy shows an amplitude larger than at 10^{14} eV and a different phase (ApJL 692, 2009, 130). Different checks of stability of the detector and consistency of the data are presented. The significance of the observation for the understanding of cosmic-ray propagation is discussed.

Presenter: Dr GHIA, Piera Luisa (LPNHE (CNRS))

Session Classification: Anisotropy measurements and methods