## Large scale anisotropy of cosmic rays and directional neutrino signals from Galactic sources

Friday, 27 September 2013 17:35 (25 minutes)

Quite recently the IceCube Collaboration has reported an observation of 26 neutrino candidates above ~ 50 TeV. Including the two ~ 1 PeV neutrinos reported earlier in 2013, these 28 events constitute a 4.3\sigma excess compared to the atmospheric background. In this talk, I will explore the compatibility between the data and an unbroken power-law neutrino spectrum, for various values of spectral index \Gamma >= 2. I will show that \Gamma ~ 2.3 is consistent at the ~ 1.5\sigma level with the observed events up to 1 PeV and to the null observation of events at higher energies. I will then assume that the sources of this unbroken spectrum are Galactic, and deduce (i) an energy-transfer fraction from parent protons to pions, and (ii) a discriminating test between the two most popular models ("dip" and "ankle") for the Galactic to extragalactic cosmic-ray transition. Future IceCube data will test the unbroken power law hypothesis, and, if the neutrino sources are Galactic, discriminate between the "dip" and "ankle" models of Galactic to extragalactic transition.

**Primary author:** Prof. ANCHORDOQUI, Luis (University of Wisconsin Milwaukee)

Presenter: Prof. ANCHORDOQUI, Luis (University of Wisconsin Milwaukee)

**Session Classification:** Large scale anisotropy of cosmic rays and directional neutrino signals from Galactic sources - Luis Anchordoqui, UW-Milwaukee

Track Classification: Luis Anchordoqui