

DECAcube – ICEcube++ - PART II

A large optical extension of IceCube for future neutrino-astronomy in the TeV-PeV range on the DecaCube-scale

David Altmann¹ and Christopher Wiebusch²



1. Estimates for in ice veto
2. Ideas for an Air-Cherenkov surface veto


2

RWTH AACHEN
IceCube





IceCube

Baseline



- We are observing a **weak astrophysical flux** with IceCube below the threshold of new technologies (Radio & Acoustics)
- **Add 100 optical strings** (invest 80 M€, 5 years installation)
- Free design parameter: **string spacing**: 120m->360m
- **Results for Tracks**
 - **Effective area** increases about linearly with spacing
 - **Energy threshold** scales moderately with spacing (2TeV (120m), 10TeV (240m), 50 TeV (360m))
⇒ galactic sources still in reach !
- **Results for Cascades**
 - Effective Volume for Cascades increases quadratically with spacing according to increase of geometrical volume
- **How about Veto and starting track capabilities?**
We have learned, these capabilities are crucial for a next generation instrument which wants to detect the sources!

Symmetrical geometry (spacing 240m)





- +100 strings
- 4-3 layers of outer strings around IceCube
- $\sim 7 \text{ km}^3$ volume

240m results:

- Muons: 3x IC3
- Cascades : 7x IC3
- $E_{\text{thresh}} \leq 10 \text{ TeV}$

Starting events estimation (A)




HESE volume
0.4 Gton

Outer strings improve veto :

- ✓ 3 outer layers more than sufficient (Result from DeepCore)
- Bottom veto is obsolete
- Dustlayer veto is obsolete
- Full IceCube volume can be used to the side - edges
- + 1 outer ring if 4 layers

Unidirectional DOMS & High QE PMT


-> better Veto !



Config	eff. Vol. Gton	#events >30TeV /a
HESE	0.4	14/a
Full IC3	0.9	31/a
+ 1 Ring	1.4-1.8	49-63 /a

DecaCube strongly improves starting event capabilities

Starting events estimation (B)




Assume:

- One outer layer veto (but thicker Top Veto)
- atm. μ veto threshold increases with spacing
- No unidirectional DOMS & High QE PMT

➤ HESE volume can be extended if threshold is raised by factor ~ 2


Note that atmospheric neutrino BG (prompt/conventional) decrease



Config	Volume Gton	#events >60TeV /a	#events >200 TeV /a	#events > 1PeV /a
HESE	0.4	8	3	1
+ 1 outer veto ring	3-5	60-100	22-37	7-12

DecaCube will yield a factor 10 increased Ernie&Bert sample


Ideal world: Surface veto




Extended IceTop (of course)

not possible to the most interesting **horizontal regions**

- Exploit (non-)imaging air cherenkov technique
- New telescope concept **FAMOUS** for FD detection
 - Based on **SiPM** 30-40% uptime (demonstrated by FACT)
 - **imaging** strongly helps in NSB background rejection (1/3000) and allows a lower threshold (critical for inclined directions)
 - VHE **gamma astronomy**
 - **Acceptance** similar to non-imaging
- **Price** \propto # pixel \propto 1/ (FOV/pixel)
- $E_{\text{thresh}} \sim 50 \text{ TeV (p)} \propto (\text{FOV})^{1/2}$




FAMOUS





7 Pixel Prototype



FOV limited by maximum SiPmt size




- **Camera pixel =**
Winston cone + 6x6 mm² SiPM
- 1.5 ° field of view per pixel
- 12 ° field of view in total






Niggemann et al ICRC 2013, 0014

Designed for Fluorescence Detection in Auger ... should work for Air Cherenkov, too


Duty Cycle of SiPM cameras: Demonstration by FACT



- Very **stable operation** (~5%) also **during full moon** demonstrated
- SouthPole: **40%/year** achievable ? (needs in.situ tests)
- SiPM **cannot** be **damaged** by bright light (PMT need protection)
- **No High voltage** needed



Showers recorded during observation of the full moon



See Knoetig et al (ICRC 2013 – 695) and Bretz et al (ICRC 2013 - 682)





First light with the FAMOUS-7 prototype



Auger Students, in the background: University hospital RWTH

FAMOUS

Energy threshold



Energy threshold : Opening window $\varnothing 55\text{cm} \rightarrow 0.25\text{m}^2$ (Hiscore: 1m^2)



FACT = $400\text{GeV} @ 9.5\text{m}^2 \Rightarrow 15\text{ TeV } (\gamma)$

Factor 2 for hadrons: $\Rightarrow 30\text{TeV}$

Factor 2-4 for horizontal hadrons \Rightarrow 60-120 TeV

- **Sub threshold data** possible, when triggered by InIce
- **Entrance window** can be enlarged

Geometrical considerations



Cover region $40^\circ < \theta < 80^\circ$
(most interesting)

Enlarged IceTop

8 km

Opening radius of LDF at
surface: r

$$\approx \tan(\theta_c) \cdot 10 \text{ km} / \cos(\theta)$$

$$\approx 1 \text{ km} / (6 \cdot \cos(\theta))$$

40° : 220m 80° : 960m

Exponentially increasing
threshold outside


40°

3 km

DecaCube

- Single Circle of air cherenkov telescopes
looking towards horizon, close to IceTop
- Field of view aligned towards IceCube

Cost estimate



6x6mm SiPMT FOV 1.6° per pixel $\rightarrow 6.12 \cdot 10^{-4}$ sr/pix
 For a zenith region $> 40^\circ - 80^\circ = 1.18\pi$ sr = 3.72 sr } $\Rightarrow \sim 6000$ pixels


Price per Telescope (Famous Prototype) 64 Pixel=256 channel		total
Pixel: 6x6 mm ² SiPm (4 chan.)	250€ x 64	16k€
Electronic (Easy Roc) 32 channel +LV	1000€ x 8	8k€
Mechanics, Lens	1200€	1.2k€
Comms + SC	1000€	1k€
Total	400€/pixel	26.2k€

*# Pixel $\propto 1/$ FOV
 FOV limited by (current)
 maximum SiPmt size*

\Rightarrow **2.4M€**

- **Costs can be greatly reduced** for mass production (10\$/piece for 100k SiPM) and larger SiPM area (FOV) possible
- Compared to 100 strings (80M€), this is a **small investment** for a **significantly improved veto @30% duty cycle**
- **Full sky coverage** can be achieved- (factor 2 of costs)
 - **Hybrid cross calibration with IceTop&IceCube**, **energy calibration of IceTop with fluorescence light**, **gamma astronomy**, **CR below the knee**

Summary




- DecaCube with 240 m spacing is a good baseline to reach $\sim 7\text{km}^3$
- Might achieve significantly improved starting events capabilities with a factor 10 increased event rates, compared to IceCube
- A surface veto based on SiPM air cherenkov detectors in addition to a conventional IceTop extension looks promising to achieve a horizontal veto with 30-40% duty cycle and $< 50\text{TeV}$ primary threshold

More work needed:

- Full MC simulation of DecaVube
 - Redo simulations with an optimized geometry
 - Test for veto capabilities
 - Simulate surface veto

Imaging versus non-imaging




- Non imaging allows a **wide field of view** with much **fewer channels** -> lower price
- **HIGHSCORE** acceptance: $A \approx 1\text{sr} \cdot 0.5\text{m}^2 = 0.5\text{ m}^2\text{ sr}$
- **FAMOUS** acceptance/pixel $A \approx 6.12 \cdot 10^{-4}\text{ sr} \cdot 0.25\text{ m}^2 = 1.5 \cdot 10^{-4}\text{ sr}$
- Night sky background (La Palma): $3.3 \cdot 10^{12}\text{ (m}^2\text{ s sr)}^{-1}$
- NSB: **Highscore/Famous**: $1,700 \cdot 10^9\text{ s}^{-1} / 0.5 \cdot 10^9\text{ s}^{-1} \approx 3000$
- **Larger SiPM** would allow for a compromise of smaller FOV (threshold) and # channels

- Need more investigations:
 - Energy threshold is critical for large inclinations
 - Boreal lights at the South Pole
 - Operation during moon

Question is open:


⇒ Need a dedicated MonteCarlo Study

Spacing 1 – 120m




Type: NuMu
E(GeV): 7.99e+06
Zen: 80.85 deg
Azi: 232.33 deg
NTrack: 11/11 shown, min E(GeV) == 12.27
NCasc: 100/2847 shown, min E(GeV) == 6.27

$E_{\nu} \approx 8$ PeV




⇒ spacing unnecessarily small !

Spacing 3: 360m




Type: NuMu
E(GeV): **3.89e+08** start-energy
Zen: 78.01 deg
Azi: 151.75 deg
NTrack: 11/11 shown, min E(GeV) == 22.21
NCasc: 100/3772 shown, min E(GeV) == 5.54




⇒ the muon is observed by a single string/layer

Spacing 3: 360m




```
Type: NuMuBar ~ 18 TeV
E(GeV) 1.87e+04
Zen: 99.36 deg
Azi: 209.08 deg
NTrack: 1/1 shown, min E(GeV) -- 16282.32
NCasc: 100/611 shown, min E(GeV) -- 1.35
```




⇒ Even for large spacing the threshold will be of the order of a few 10 TeV (energy loss increases linearly)

Spacing 2





Type: NuMuBar
E(GeV): **4.42e+07** ~ 40 PeV
Zen: 78.72 deg
Azi: 141.19 deg
NTrack: 11/11 shown, min E(GeV) == 5.45
NCasc: 100/4250 shown, min E(GeV) == 1.30



⇒ In DecaCube-2 the muon is observed by several strings/layer



Improvement Factor w.r. IceCube-86



Cascades (new result) Improvement Factor w.r. IceCube-86



- Events contained
- Nchannel ≥ 15
- Nstring ≥ 3

Glashow resonance

- Improvement \propto Volume $\propto d^2$
- Large spacing fully effective only > 30 PeV

**Go for now
with option 2**