

Search for point-like sources in the astrophysical muon neutrino flux with IceCube

Monday, 8 May 2017 17:30 (15 minutes)

Using a sample of neutrino-induced muon track events, IceCube has confirmed the high-energy diffuse astrophysical neutrinos flux, first found in a sample of high energy starting events. In order to identify the sources of this diffuse flux, these muon-neutrino events are ideal because of their excellent angular resolution. Here we present a search for point-like neutrino sources based on the same sample that was used to measure the diffuse high-energy astrophysical muon-neutrino flux with a livetime of seven years. This high purity sample is restricted to the Northern hemisphere and consists of ~425000 events with a median angular resolution of $\sim 1^\circ$ at 1 TeV that decreases to $\sim 0.3^\circ$ at 1 PeV. In this analysis an unbinned likelihood maximization is used that is optimized for point-like neutrino emission with the same characteristics as the diffuse muon neutrino flux. The sensitivity improves for sources with an E^{-2} spectrum by 30% compared to previous analysis and is at a level of $E^2 \partial\phi/\partial E = 3 \cdot 10^{-13} \text{ TeV cm}^{-2} \text{ s}^{-1}$. We report about the status of this search and present an analysis searching for a population of neutrino sources, which are too weak to be significant individually.

Primary author: Mr REIMANN, René (RWTH Aachen)

Presenter: Mr REIMANN, René (RWTH Aachen)

Session Classification: Neutrino Astronomy

Track Classification: Neutrino Astronomy - Convenor: Gisela Anton, FAU / ECAP